



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

given by

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = v^{v/(v-1)} \quad \text{and} \quad \frac{z^2}{c^2} = v^{1/(v-1)} - 1.$$

This surface is symmetrical with reference to the plane $z=0$, and incloses that plane by an infinite elliptic boundary. As v increases from 0 to ∞ the sections parallel to $z=0$ decrease continually, remaining ellipses always, until they reach their limiting size, which is that of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$.

Another companion to the same curve is

$$\frac{x^2}{a^2} - \frac{z^2}{c^2} = v^{v/(v-1)} \quad \text{and} \quad \frac{y^2}{b^2} = 1 - v^{1/(v-1)}.$$

y has a real value only when $v^{1/(v-1)} = 1$. Hence the companion degenerates to $y=0$ and $\frac{x^2}{a^2} - \frac{z^2}{c^2} = \infty$. It is therefore an hyperbola infinitely removed from the origin and lying in the plane $y=0$.

The companion to the sphere $x^2 + y^2 + z^2 = a^2$ is

$$x^2 + y^2 = v^{v/(v-1)} \quad \text{and} \quad z^2 = a^2 - v^{1/(v-1)}.$$

When $a^2=1$, the companion is an infinite circle. When $a^2 < 1$, the companion is imaginary. When $a^2 > 1$, the companion is a real surface. Every section by planes parallel to $z=0$ is a circle. For all values of v which make $v^{1/(v-1)} > a^2$ the planes which make circular sections are imaginary.



NOTE ON THE ADDITION THEOREM IN TRIGONOMETRY.

By DR. G. A. MILLER.

When $\cos \alpha$, $\cos \beta$, $\sin \alpha$, $\sin \beta$ are substituted for x_1 , x_2 , y_1 , y_2 respectively, in the well known identity

$$(x_1^2 + y_1^2)(x_2^2 + y_2^2) = (x_1 y_2 + x_2 y_1)^2 + (x_1 x_2 - y_1 y_2)^2,$$

there results

$$(\cos^2 \alpha + \sin^2 \alpha)(\cos^2 \beta + \sin^2 \beta) = (\cos \alpha \sin \beta + \cos \beta \sin \alpha)^2 + (\cos \alpha \cos \beta - \sin \alpha \sin \beta)^2$$

or

$$(\sin \alpha \cos \beta + \cos \alpha \sin \beta)^2 + (\cos \alpha \cos \beta - \sin \alpha \sin \beta)^2 = 1.$$

For every value of α and β the two expressions in parenthesis represent real

numbers whose squares are together equal to unity. Hence one represents $\sin\phi$ and the other $\cos\phi$, where ϕ is a function of α and β .

Suppose we know that

$$\sin\alpha \cos\beta + \cos\alpha \sin\beta = \sin(\alpha + \beta) = \sin(180^\circ - \alpha - \beta) \quad (A).$$

It follows from the given identity that

$$\cos\alpha \cos\beta - \sin\alpha \sin\beta = \cos(\alpha + \beta) \text{ or } \cos(180^\circ - \alpha - \beta).$$

The latter is impossible since $\alpha = \beta = 0$ would not satisfy the equation. Hence we have that from the given identity and (A) it follows that

$$\cos\alpha \cos\beta - \sin\alpha \sin\beta = \cos(\alpha + \beta) = \cos(-\alpha - \beta).$$

If the given identity is written in the form

$$(x_1^2 + y_1^2)(x_2^2 + y_2^2) = (x_2y_1 - x_1y_2)^2 + (x_1x_2 + y_1y_2)^2$$

we have, by the method used above, that

$$\sin\alpha \cos\beta - \cos\alpha \sin\beta \text{ and } \cos\alpha \cos\beta + \sin\alpha \sin\beta$$

are respectively the sign and cosine of a given angle. It should be observed that it does not matter which of these expressions is taken for $\sin\phi$ except that the formulas come out in a little briefer form if we proceed as above.

This note does not aim at completeness. Its object is to call the attention of teachers of trigonometry to the fact that the derivation of these formulas could proceed along lines which differ from those which are generally pursued, and that these methods offer many advantages. Other things being equal it is always desirable to exhibit connection between different formulas rather than to derive each independently.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

229. Proposed by BENJAMIN FRANKLIN YANNEY, Mount Union College, Alliance, Ohio.

If $a_1^n + a_2^n + a_3^n + \dots + a_r^n = A^n$, $a_1^m + a_2^m + a_3^m + \dots + a_r^m >$ or $< A^m$, according as $m <$ or $> n$; provided all the letters stand for positive real numbers.